

CXL at the Slit Lamp: No Clinically Relevant Changes in Corneal Riboflavin Distribution During Upright UV Irradiation

To the Editor:

Photo-activated chromophore for keratitis cross-linking (PACK-CXL)¹⁻⁴ may become an attractive treatment for developing and emerging countries where access to ophthalmic care and medical equipment is limited.⁵ PACK-CXL treatment has the potential to be performed at the slit-lamp within 3 to 5 minutes.³ We have addressed a remaining concern: is the stromal riboflavin distribution affected by gravity if the ultraviolet irradiation takes place in the upright position?

Freshly enucleated porcine eyes were incubated in 0.1% hypotonic riboflavin solution without dextran for 30 minutes and then divided into three groups: corneas fixed in a vertical position for 30 minutes (n = 4), corneas fixed in a vertical position for 60 minutes (n = 8), and corneas fixed in a horizontal position for 60 minutes (n = 8). Four corneas were immersed in 0.9% sodium chloride for 60 minutes and served as fluorescence controls. The fluorescence signal of each cornea was measured with a spectrophotometer. Statistical significance was determined with Student's *t* tests and a confidence interval of 95%.

After 30 minutes, there was no difference ($P = .22$) between superior and inferior riboflavin saturation, whereas a statistically significant ($P = .002$) gradient of 3.36% was found after 60 minutes (**Figure 1**).

Gravitational influence on the riboflavin distribution was observed only after 60 minutes of vertical positioning. Given that a PACK-CXL treatment typically lasts for 3 to 30 minutes,^{2,4} this difference can be considered not clinically relevant. Limitations of this study include a small sample size and the use of porcine instead of human corneas.

REFERENCES

1. Hafezi F, Randleman JB. PACK-CXL: defining CXL for infectious keratitis. *J Refract Surg.* 2014;30:438-439.




Figure 1. Changes (%) in fluorescence levels along the vertical axis in riboflavin-soaked corneas mounted in a vertical position for 30 and 60 minutes, respectively.

2. Iseli HP, Thiel MA, Hafezi F, Kampmeier J, Seiler T. Ultraviolet A/riboflavin corneal cross-linking for infectious keratitis associated with corneal melts. *Cornea.* 2008;27:590-594.
3. Richoz O, Kling S, Hoogewoud F, et al. Antibacterial efficacy of accelerated photoactivated chromophore for keratitis-corneal collagen cross-linking (PACK-CXL). *J Refract Surg.* 2014;30:850-854.
4. Said DG, Elalfy MS, Gatziofias Z, et al. Collagen cross-linking with photoactivated riboflavin (PACK-CXL) for the treatment of advanced infectious keratitis with corneal melting. *Ophthalmology.* 2014;121:1377-1382.
5. Whitcher JP, Srinivasan M. Corneal ulceration in the developing world: a silent epidemic. *Br J Ophthalmol.* 1997;81:622-623.

Basile Salmon
Olivier Richoz, MD, PhD
David Tabibian, MD
Sabine Kling, PhD
Raphael Wuarin
Farhad Hafezi, MD, PhD
 Geneva, Switzerland

Drs. Richoz and Hafezi are co-inventors of the PCT/CH2012/000090 application (UV irradiation device), and Dr. Hafezi is Chief Scientific Officer and shareholder of EMAGine. The remaining authors have no financial or proprietary interest in the materials presented herein.

doi:10.3928/1081597X-20161219-03